



Northeast Fisheries Science Center Reference Document 09-07

Determination of Conversion Factors for Vessel Comparison Studies

by Henry O. Milliken and Michael J. Fogarty

April 2009

Recent Issues in This Series

- 08-09 *Bycatch of Harbor Porpoises in Three U.S. Gillnet Management Areas: Southern Mid-Atlantic, Offshore, and Western Gulf of Maine*, by CD Orphanides and DL Palka. July 2008.
- 08-10 *Harbor Porpoise Bycatch Rates that Indicate Compliance with Pinger Regulations for the Northeast Gillnet Fishery*, by DL Palka and CD Orphanides. July 2008.
- 08-11 *47th SAW Assessment Summary Report*, by the 47th Northeast Regional Stock Assessment Workshop (47th SAW). July 2008.
- 08-12 *47th SAW (a) Assessment Report and (b) Appendixes*, by the 47th Northeast Regional Stock Assessment Workshop (47th SAW). July 2008.
- 08-13 *Predicted Habor Porpoise Bycatch under Potential Mitigation Measure Scenarios*, by DL Palka and CD Orphanides. August 2008.
- 08-14 *Predicted Bycatch of Harbor Porpoises under Various Alternatives to Reduce Bycatch in the US Northeast and Mid-Atlantic Gillnet Fisheries*, by DL Palka and CD Orphanides. In press.
- 08-15 *Assessment of 19 Northeast Groundfish Stocks through 2007: A Report of the 3rd Groundfish Assessment Review Meeting (GARM III)*, Northeast Fisheries Science Center, Woods Hole, Massachusetts, August 4-8, 2008, by Northeast Fisheries Science Center. August 2008.
- 08-16 *Assessment of 19 Northeast Groundfish Stocks through 2007: A Report of the 3rd Groundfish Assessment Review Meeting (GARM III) -- Appendixes*, by Northeast Fisheries Science Center. September 2008.
- 08-17 *Preparation of the Northeast Fisheries Observer Program Gillnet Data for Use in Bycatch Analyses of Protected Species*, by ML Warden and CD Orphanides. August 2008.
- 08-18 *A Description of the Allocation Procedure Applied to the 1994 to 2007 Commercial Landings data*, by SE Wigley, P Hersey, and JE Palmer. September 2008.
- 08-19 *11th Flatfish Biology Conference Program and Abstracts, Dec. 3-4, 2008, Water's Edge Resort and Spa, Westbrook, Connecticut*, by Conference Steering Committee: R Mercaldo-Allen (Chair), A Calabrese, D Danila, M Dixon, A Jearld, T Munroe, Deborah Pacileo, C Powell, and S Sutherland. November 2008.
- 08-20 *Estimated average annual bycatch of loggerhead sea turtles (Caretta caretta) in US Mid-Atlantic bottom otter trawl gear, 1996-2004 (2nd edition)*, by KT Murray. November 2008.
- 09-01 *Report of the Retrospective Working Group, January 14-16, 2008, Woods Hole, Massachusetts*, by CM Legault, Chair. January 2009.
- 09-02 *The Northeast Data Poor Stocks Working Group Report, December 8-12, 2008 Meeting*, by Northeast Data Poor Stocks Working Group. January 2009.
- 09-03 *The 2008 Assessment of the Gulf of Maine Atlantic Cod (Gadus morhua) Stock*, by RK Mayo, G Shepherd, L O'Brien, LA Col, and M. Traver. February 2009.
- 09-04 *Mortality and serious injury determinations for baleen whale stocks along the United States eastern seaboard and adjacent Canadian maritimes, 2003-2007*, by AH Glass, TVN Cole, and M Garron. March 2009.
- 09-05 *North Atlantic Right Whale Sighting Survey (NARWSS) and Right Whale Sighting Advisory System (RWSAS) 2008 Results Summary*, by C Khan, TVN Cole, P Duley, AH Glass, M Niemeyer, and C Christman. March 2009.
- 09-06 *A Bibliography of the Long-Finned Pilot Whale, Globicephala melas, and the Short-Finned Pilot Whale, Globicephala macrorhynchus, in the North Atlantic Ocean*, compiled by FW Wenzel, JR Nicolas, A Abend, and B Hayward. April 2009.

Determination of Conversion Factors for Vessel Comparison Studies

by Henry O. Milliken and Michael J. Fogarty

Northeast Fisheries Science Center, 166 Water Street, Woods Hole, MA 02543

U.S. DEPARTMENT OF COMMERCE
National Oceanic and Atmospheric Administration
National Marine Fisheries Service
Northeast Fisheries Science Center
Woods Hole, Massachusetts

April 2009

Northeast Fisheries Science Center Reference Documents

This series is a secondary scientific series designed to assure the long-term documentation and to enable the timely transmission of research results by Center and/or non-Center researchers, where such results bear upon the research mission of the Center (see the outside back cover for the mission statement). These documents receive internal scientific review, and most receive copy editing. The National Marine Fisheries Service does not endorse any proprietary material, process, or product mentioned in these documents.

All documents issued in this series since April 2001, and several documents issued prior to that date, have been copublished in both paper and electronic versions. To access the electronic version of a document in this series, go to <http://www.nefsc.noaa.gov/nefsc/publications/>. The electronic version is available in PDF format to permit printing of a paper copy directly from the Internet. If you do not have Internet access, or if a desired document is one of the pre-April 2001 documents available only in the paper version, you can obtain a paper copy by contacting the senior Center author of the desired document. Refer to the title page of the document for the senior Center author's name and mailing address. If there is no Center author, or if there is corporate (*i.e.*, non-individualized) authorship, then contact the Center's Woods Hole Laboratory Library (166 Water St., Woods Hole, MA 02543-1026).

This document's publication history is as follows: manuscript submitted for review March 23, 2009; manuscript accepted through technical review April 27, 2009; manuscript accepted through policy review April 27, 2009; and final copy submitted for publication April 27, 2009. Pursuant to section 515 of Public Law 106-554 (the Information Quality Act), this information product has undergone a pre-dissemination review by the Northeast Fisheries Science Center, completed on April 27, 2009. The signed pre-dissemination review and documentation is on file at the NEFSC Editorial Office. This document may be cited as:

Milliken HO, Fogarty MJ. 2009. Determination of Conversion Factors for Vessel Comparison Studies. US Dept Commer, Northeast Fish Sci Cent Ref Doc. 09-07; 18 p. Available from: National Marine Fisheries Service, 166 Water Street, Woods Hole, MA 02543-1026, or online at <http://www.nefsc.noaa.gov/nefsc/publications/>

Table of Contents

ABSTRACT	iv
INTRODUCTION	1
MATERIALS & METHODS	1
RESULTS	2
DISCUSSION	3
ACKNOWLEDGEMENTS	3
LITERATURE CITED	3
APPENDIX A	9

List of Tables

Table 1. Cruise information for paired fishing power cruises between the R/V's <i>Albatross IV</i> and <i>Delaware II</i> since 1982	4
Table 2. A list of the species analyzed, the ratio estimates (conversion factors), the standard error of the estimates, and 90% confidence limits	4

List of Figures

Figure 1. Station locations for paired comparisons	5
Figure 2. Confidence intervals for "GLM" conversion factors versus the "ratio" conversion factors for the 1980s data only	6
Figure 3. A comparison of the ratio conversion factors derived from the 1980s data and the data following the refit of the R/V <i>Delaware II</i>	7
Figure 4. Plot of the ratio between the sum of the R/V <i>Albatross IV</i> divided by the sum of the R/V <i>Delaware II</i> and the 90% confidence intervals	8

ABSTRACT

NOAA's Northeast Fisheries Science Center has conducted groundfish surveys on the Northeast Continental Shelf of the United States in autumn since 1963. These surveys are used to compute relative abundance indices of fish and selected invertebrates. Since the inception of the survey program, the R/V *Albatross IV* has been the primary vessel conducting the work, except periodically when the R/V *Delaware II* has been utilized because the R/V *Albatross IV* was not available. When the R/V *Delaware II* was used, differences in vessel characteristics necessitate the development of conversion coefficients to adjust catch rates using the R/V *Albatross* as the standard.

A series of vessel calibration studies between the R/Vs *Albatross IV* and *Delaware II* employing a paired tow design were initiated in 1981. In 1997, the R/V *Delaware II* underwent a major refit. Accordingly, for these two research vessels, there are two variations in the experimental time series that require consideration in development of calibration coefficients. We explored the use of a ratio estimator for the development of a vessel conversion coefficient and compared these with a general linear model (GLM) approach used by Byrne and Forrester (1991).

INTRODUCTION

For many decades the Federal Government has conducted groundfish surveys out of Woods Hole, MA. With the arrival of the newly built research vessel R/V *Albatross IV* in 1963, the Bureau of Commercial Fisheries (now National Marine Fisheries Service) began a time series that serves as the basis for some of the longest time series of standardized fishery-independent indices of relative abundance in the world. In addition to tracking abundance of mature animals, research surveys provide indices of juvenile abundance, which can indicate strong year classes before fish are vulnerable to commercial or recreational fisheries. Research surveys indicate the status of a stock over its entire range, not just in small areas of commercial or recreational concentration. Surveys also provide data on growth, maturity, predation, and mortality of a stock as well as trophic dynamics of fish communities (Azarovitz 1981).

The R/V *Albatross IV*'s autumn bottom trawl survey was conducted using a #36 Yankee net with a sixty-foot (18.3 m) headrope and eighty-foot (24.4 m) footrope. The sweep was comprised of sixteen-inch (40.6 cm) rollers that allow the gear to be fished on most bottom types. The net and fishing operation remained almost the same since the start of the autumn survey in 1963 (Azarovitz 1981). When changes were made, comparative tows were performed to quantify the changes in the catchability of the gear. Once determined, a conversion coefficient was applied to the data so that any differences in the catchability of the gear could be compared to future catches (Fogarty 1997).

Conversion coefficients between the R/V's *Albatross IV* and *Delaware II* were developed by Byrne and Forrester (1991) using a general linear model (GLM) approach. When the R/V *Delaware II* was used for surveys, these estimates were used to transform catch data. In 1997 the R/V *Delaware II* underwent a major refit. This refit included changing the winches from slower direct drive to faster free spooling winches. After the refit, the previous conversion factors were still employed to transform the data. However, the GLM approach does not allow the consideration of tows with zero catch, thus those data were lost. An alternative approach using ratio estimators, which use a ratio of the sum of the paired data, allows the comparison of all catch data between the two vessels.

Because the GLM method does not permit the inclusion of paired tows where one of the tows has zero catch, we compared the use of the GLM and ratio estimator methods for deriving the conversion factors. Additionally we examined the conversion factors before and after the refit of the R/V *Delaware II* in 1997.

MATERIALS & METHODS

The R/V's *Albatross IV* and *Delaware II* fished alongside each other five times since the R/V *Delaware II* was refurbished in 1997 (Table 1, Figure 1). During these cruises, the R/V *Delaware II* accompanied the R/V *Albatross IV* while the R/V *Albatross IV* was engaged in its standard bottom trawl survey. Three paired comparison cruises from the 1980s were used to compare the GLM and ratio estimators for developing conversion factors. One 1998 cruise and four cruises from the 2000s were utilized to construct the ratio estimator model to estimate conversion coefficients.

The conversion coefficient (α) was calculated as:

$$\hat{\alpha} = \frac{\sum_{i=1}^n w V_{2_i} V_{1_i}}{\sum_{i=1}^n w V_{1_i}^2}$$

where w is a weighting coefficient that depends on the relationship between the variance of V2 (Vessel 2) and V1(Vessel 1); the variance is defined as:

$$\hat{\text{var}}\left(\hat{\alpha}\right) = \frac{\frac{1}{n-1} \sum_{i=1}^n w \left(V_{2_i} - \hat{\alpha} V_{1_i}\right)^2}{\sum_{i=1}^n w V_{1_i}^2}$$

If the variance of V2 and V1 is proportional, the weighting coefficient is V1-1 and the estimator for the conversion coefficient is:

$$\hat{\alpha} = \frac{\sum_{i=1}^n V_{2_i}}{\sum_{i=1}^n V_{1_i}}$$

We used the proportional variance estimator in our analyses where “V2” is the variance of the R/V *Albatross IV* and “V1” is the variance of the R/V *Delaware II*.

Conversion coefficients were developed for winter skate (*Leucoraja ocellata*), silver hake (*Merluccius bilinearis*), Atlantic cod (*Gadus morhua*), haddock (*Melanogrammus aeglefinus*), white hake (*Urophycis tenuis*), red hake (*Urophycis chuss*), American plaice (*Hippoglossoides platessoides*), fourspot flounder (*Hippoglossina oblonga*), yellowtail flounder (*Limanda ferruginea*), witch flounder (*Glyptocephalus cynoglossus*), windowpane flounder (*Scophthalmus aquosus*), redfish (*Sebastes fasciatus*), and ocean pout (*Zoarces americanus*).

RESULTS

When we examined the conversion factors derived from both the GLM and ratio estimator models using the 1980s data, with the exception of haddock, there appears to be good correlation between the two methods for determining vessel conversion coefficients (Figure 2).

Ratio conversion coefficients calculated for catches before and after the R/V *Delaware*'s 1997 refit suggest that the vessel effect was reduced after the vessel was modified. Twelve of the 14 species' coefficients calculated from the 1980s data showed a difference, while only 4 of the 14 species show a difference after the refit (Figure 3). The species that showed a difference included cod, haddock, pollock and redfish (Figure 4). For these four species, the post-fit

conversion factors should be applied to the R/V *Delaware II* data when the R/V *Delaware II* was used in place of the R/V *Albatross IV* for the standardized groundfish surveys (Table 2).

DISCUSSION

Differences in catch efficiencies between the R/V *Albatross IV* and R/V *Delaware II* were reduced after the refit of the R/V *Delaware II*. It was unanticipated that the refit would reduce the difference in the catchability between the two vessels or, in other words, reduce the catchability of the R/V *Delaware II*. After the R/V *Delaware II* was modified, the winches were replaced with faster winches. We hypothesize that because the winches were faster, the amount of time the net spent on the bottom was reduced. The winch speed was designed to match the speed of the winches on the R/V *Albatross IV* which appears to have equalized the catchability between the two vessels.

The ratio conversion coefficients were calculated for fourteen species. The next step will be to apply it to the other principal groundfish species to determine if conversion coefficients are required.

ACKNOWLEDGEMENTS

This work would not be possible without the efforts of numerous members of the Northeast Fisheries Science Center, particularly the Resource Survey Branch, who participated in these cruises and helped collect this data. A special thanks is due to Kathy Mays of the Northeast Fisheries Science Center who spent considerable time helping to develop the proper scripts to pull the data from the survey database.

LITERATURE CITED

- Azarovitz TR. 1981. A brief historical review of the Woods Hole Laboratory trawl survey time series. *Can Spec Pub Fish Aq Sci.* 58:62-67.
- Byrne CJ, Forrester JRS. 1991. Relative Fishing Power of NOAA R/V's *Albatross IV* and *Delaware II*. In: Report of the Twelfth Northeast Regional Stock Assessment Workshop. US. Dept. Commer., NOAA, Northeast Fisheries Science Center Ref. Doc. 91-03, 187 p.
- Fogarty M.J. 1997. Standardizing fishing power in research vessel surveys. In: Proceedings of the Workshop on Maintaining Current & Future Fisheries Resource Survey Capabilities. Special Report No. 63 of the Atlantic States Marine Fisheries Commission, p 54-63.

Table 1. Cruise information for paired fishing power cruises between the R/V's *Albatross IV* and *Delaware II* since 1982. When the start date for the *Albatross IV* precedes the *Delaware II* it is because the comparative work did not occur during all four legs of a survey.

CRUISE NUMBER	VESSEL NAME	START DATE	END DATE	SURVEY	NUMBER OF PAIRS	GEAR TYPE
198206	Albatross IV	9/13/1982	11/12/1982	FALL	144	Yankee 36
198207	Delaware II	10/4/1982	10/29/1982			
198306	Albatross IV	9/12/1983	11/10/1983	FALL	113	Yankee 36
198307	Delaware II	10/4/1983	10/26/1983			
198803	Albatross IV	9/12/1988	10/28/1988	FALL	227	Yankee 36
198804	Delaware II	9/19/1988	10/28/1988			
199802	Albatross IV	3/2/1998	4/20/1998	SPRING	44	Yankee 36
199807	Delaware II	3/30/1998	4/10/1998			
200002	Albatross IV	3/16/2000	5/3/2000	SPRING	125	Yankee 36
200003	Delaware II	3/20/2000	4/14/2000			
200102	Albatross IV	2/26/2001	4/30/2001	SPRING	69	Yankee 36
200104	Delaware II	3/26/2001	4/20/2001			
200202	Albatross IV	3/4/2002	4/26/2002	SPRING	131	Yankee 36
200203	Delaware II	4/3/2002	4/26/2002			
200209	Albatross IV	9/3/2002	10/25/2002	FALL	51	Yankee 36
200211	Delaware II	10/16/2002	10/25/2002			

Table 2. A list of the species analyzed, the ratio estimates (conversion factors), the standard error of the estimates, and 90% confidence limits.

SPECIES	RATIO ESTIMATE	STANDARD ERROR	90% CONFIDENCE INTERVALS	
Winter Skate	0.93	0.06	0.831	1.029
Silver Hake	1.02	0.07	0.9045	1.1355
Cod	0.82	0.06	0.721	0.919
Haddock	0.83	0.06	0.731	0.929
Pollock	0.69	0.05	0.6075	0.7725
White Hake	1.04	0.07	0.9245	1.1555
Red Hake	0.91	0.06	0.811	1.009
American Plaice	1.01	0.07	0.8945	1.1255
Fourspot Flounder	1.04	0.07	0.9245	1.1555
Yellowtail Flounder	1.07	0.07	0.9545	1.1855
Witch Flounder	1.00	0.07	0.8845	1.1155
Windowpane Flounder	0.95	0.07	0.8345	1.0655
Redfish	0.87	0.06	0.771	0.969
Ocean Pout	0.98	0.07	0.8645	1.0955




Figure 1. Station locations for paired comparisons (listed in Table 1). Different colors signify different cruises.




Figure 2. Confidence intervals (95%) for “GLM” conversion factors versus the “ratio” conversion factors for the 1980s data only. Species which did not have a conversion factor in one of the data sets were omitted. When the error bar falls above or below “1”, the catch weights between the two vessels are significantly different. When the bars fall below “1”, it indicates that the R/V *Delaware* caught significantly more of that species.




Figure 3. A comparison of the ratio conversion factors derived from the 1980's data (A) and the data following the refit of the R/V *Delaware II* (B). When the error bar falls above or below “1”, the catch weights between the two vessels are significantly different. When the bars fall below “1”, it indicates that the R/V *Delaware* caught significantly more of that species. Bars represent the 90% confidence intervals.




Figure 4. Plot of the ratio between the sum of the R/V *Albatross IV* divided by the sum of the R/V *Delaware II* and the 90% confidence intervals. When the error bar falls above or below “1”, the catch weights between the two vessels are significantly different. When the bars fall below “1”, it indicates that the R/V *Delaware* caught significantly more of that species.

APPENDIX A

Winter skate [cruise = Pooled data]

Dep Var: WINTERSK_ALB N: 482 Multiple R: 0.728 Squared multiple R: 0.530

Adjusted squared multiple R: 0.529 Standard error of estimate: 40.948

Effect	Coefficient	Std Error	Std Coef	Tolerance	t	P(2 Tail)
CONSTANT	4.502	1.934	0.000	.	2.328	0.020
WINTERSK_DEL	0.614	0.026	0.728	1.000	23.276	0.000


Analysis of Variance

Source	Sum-of-Squares	df	Mean-Square	F-ratio	P
Regression	908363.493	1	908363.493	541.750	0.000
Residual	804825.997	480	1676.721		


Case	83 is an outlier	(Studentized Residual =	5.259)
Case	198 has large leverage	(Leverage =	0.318)
Case	198 is an outlier	(Studentized Residual =	-15.989)
Case	198 has large influence	(Cook distance =	38.888)
Case	207 is an outlier	(Studentized Residual =	12.938)
Case	212 has large leverage	(Leverage =	0.295)
Case	216 is an outlier	(Studentized Residual =	8.753)
Case	224 has large leverage	(Leverage =	0.054)
Case	224 is an outlier	(Studentized Residual =	9.839)
Case	257 has large leverage	(Leverage =	0.068)

Durbin-Watson D Statistic 1.866
First Order Autocorrelation 0.067

Plot of Residuals against Predicted Values



Winter skate weights (kg)



Silver Hake [cruise = Pooled data]

Dep Var: SH_ALB N: 482 Multiple R: 0.343 Squared multiple R: 0.118

Adjusted squared multiple R: 0.116 Standard error of estimate: 22.614

Effect	Coefficient	Std Error	Std Coef	Tolerance	t	P(2 Tail)
CONSTANT	1.921	1.073	0.000	.	1.791	0.074
SH_DEL	0.613	0.077	0.343	1.000	8.012	0.000


Analysis of Variance

Source	Sum-of-Squares	df	Mean-Square	F-ratio	P
Regression	32832.852	1	32832.852	64.200	0.000
Residual	245479.063	480	511.415		


```
-----
Case          95 is an outlier      (Studentized Residual =    103.250)
Case          229 has large leverage (Leverage =                0.044)
Case          232 has large leverage (Leverage =                0.303)
Case          248 has large leverage (Leverage =                0.309)
Case          344 has large leverage (Leverage =                0.078)
```

Durbin-Watson D Statistic 2.011
First Order Autocorrelation -0.006

Plot of Residuals against Predicted Values



Silver hake weights (kg)



Atlantic Cod [cruise = Pooled data]

Dep Var: COD_ALB N: 482 Multiple R: 0.634 Squared multiple R: 0.401

Adjusted squared multiple R: 0.400 Standard error of estimate: 9.876

Effect	Coefficient	Std Error	Std Coef	Tolerance	t	P(2 Tail)
CONSTANT	0.969	0.470	0.000	.	2.063	0.040
COD_DEL	0.513	0.029	0.634	1.000	17.943	0.000


Analysis of Variance

Source	Sum-of-Squares	df	Mean-Square	F-ratio	P
Regression	31398.690	1	31398.690	321.946	0.000
Residual	46813.349	480	97.528		


Case	98 is an outlier	(Studentized Residual =	6.186)
Case	186 has large leverage	(Leverage =	0.053)
Case	222 has large leverage	(Leverage =	0.046)
Case	223 has large leverage	(Leverage =	0.040)
Case	236 has large leverage	(Leverage =	0.182)
Case	236 is an outlier	(Studentized Residual =	-7.502)
Case	318 has large leverage	(Leverage =	0.040)
Case	318 is an outlier	(Studentized Residual =	5.292)
Case	329 is an outlier	(Studentized Residual =	4.880)
Case	342 has large leverage	(Leverage =	0.177)
Case	359 has large leverage	(Leverage =	0.063)
Case	361 has large leverage	(Leverage =	0.051)
Case	362 is an outlier	(Studentized Residual =	17.078)
Case	383 is an outlier	(Studentized Residual =	6.324)
Case	393 has large leverage	(Leverage =	0.105)

Durbin-Watson D Statistic 1.923
First Order Autocorrelation 0.037

Plot of Residuals against Predicted Values



Atlantic cod weights (kg)



Haddock [cruise = Pooled data]

Dep Var: HADDOCK_ALB N: 482 Multiple R: 0.646 Squared multiple R: 0.417

Adjusted squared multiple R: 0.416 Standard error of estimate: 11.751

Effect	Coefficient	Std Error	Std Coef	Tolerance	t	P(2 Tail)
CONSTANT	0.619	0.550	0.000	.	1.126	0.261
HADDOCK_DEL	0.869	0.047	0.646	1.000	18.545	0.000


Analysis of Variance

Source	Sum-of-Squares	df	Mean-Square	F-ratio	P
Regression	47488.890	1	47488.890	343.901	0.000
Residual	66282.680	480	138.089		


Case	236 has large leverage	(Leverage =	0.189)
Case	325 has large leverage	(Leverage =	0.088)
Case	325 is an outlier	(Studentized Residual =	-4.370)
Case	329 is an outlier	(Studentized Residual =	11.928)
Case	340 is an outlier	(Studentized Residual =	27.927)
Case	342 has large leverage	(Leverage =	0.056)
Case	382 has large leverage	(Leverage =	0.177)
Case	383 has large leverage	(Leverage =	0.069)
Case	389 is an outlier	(Studentized Residual =	4.106)
Case	393 has large leverage	(Leverage =	0.214)

Durbin-Watson D Statistic 1.996
First Order Autocorrelation 0.002

Plot of Residuals against Predicted Values



Haddock weights (kg)



White Hake [cruise = Pooled data]

Dep Var: W_HAKE_ALB N: 482 Multiple R: 0.693 Squared multiple R: 0.480

Adjusted squared multiple R: 0.479 Standard error of estimate: 4.411

Effect	Coefficient	Std Error	Std Coef	Tolerance	t	P(2 Tail)
CONSTANT	0.621	0.209	0.000	.	2.970	0.003
W_HAKE_DEL	0.518	0.025	0.693	1.000	21.036	0.000


Analysis of Variance

Source	Sum-of-Squares	df	Mean-Square	F-ratio	P
Regression	8608.851	1	8608.851	442.493	0.000
Residual	9338.566	480	19.455		


Case	232 has large leverage	(Leverage =	0.064)
Case	233 has large leverage	(Leverage =	0.188)
Case	235 has large leverage	(Leverage =	0.059)
Case	243 is an outlier	(Studentized Residual =	4.096)
Case	252 has large leverage	(Leverage =	0.139)
Case	252 is an outlier	(Studentized Residual =	4.065)
Case	368 is an outlier	(Studentized Residual =	5.060)
Case	371 is an outlier	(Studentized Residual =	6.549)
Case	372 has large leverage	(Leverage =	0.081)
Case	372 is an outlier	(Studentized Residual =	5.901)
Case	392 has large leverage	(Leverage =	0.061)
Case	397 is an outlier	(Studentized Residual =	14.868)
Case	398 has large leverage	(Leverage =	0.094)
Case	398 is an outlier	(Studentized Residual =	-6.746)
Case	405 is an outlier	(Studentized Residual =	4.078)

Durbin-Watson D Statistic 1.994
First Order Autocorrelation 0.003

Plot of Residuals against Predicted Values



White hake weights (kg)



Red Hake [cruise = Pooled data]

Dep Var: RED_HAKE_ALB N: 482 Multiple R: 0.743 Squared multiple R: 0.552

Adjusted squared multiple R: 0.551 Standard error of estimate: 6.582

Effect	Coefficient	Std Error	Std Coef Tolerance	t	P(2 Tail)
CONSTANT	0.561	0.312	0.000	1.800	0.073
RED_HAKE_DEL	0.705	0.029	0.743	24.322	0.000


Analysis of Variance

Source	Sum-of-Squares	df	Mean-Square	F-ratio	P
Regression	25627.542	1	25627.542	591.546	0.000
Residual	20795.050	480	43.323		


Case	26 has large leverage	(Leverage =	0.060)
Case	27 has large leverage	(Leverage =	0.048)
Case	45 has large leverage	(Leverage =	0.146)
Case	45 is an outlier	(Studentized Residual =	-10.750)
Case	86 is an outlier	(Studentized Residual =	5.572)
Case	95 is an outlier	(Studentized Residual =	7.378)
Case	145 has large leverage	(Leverage =	0.085)
Case	145 is an outlier	(Studentized Residual =	11.807)
Case	155 has large leverage	(Leverage =	0.166)
Case	157 has large leverage	(Leverage =	0.136)
Case	157 is an outlier	(Studentized Residual =	5.016)
Case	258 is an outlier	(Studentized Residual =	6.661)
Case	259 is an outlier	(Studentized Residual =	4.167)
Case	287 is an outlier	(Studentized Residual =	4.699)
Case	341 has large leverage	(Leverage =	0.072)
Case	341 is an outlier	(Studentized Residual =	-6.482)
Case	346 is an outlier	(Studentized Residual =	4.290)
Case	347 has large leverage	(Leverage =	0.063)

Durbin-Watson D Statistic 1.839
First Order Autocorrelation 0.080

Plot of Residuals against Predicted Values



Red hake weights (kg)



American Plaice [cruise = Pooled data]

Dep Var: A_PLAICE_ALB N: 482 Multiple R: 0.914 Squared multiple R: 0.836

Adjusted squared multiple R: 0.835 Standard error of estimate: 1.373

Effect	Coefficient	Std Error	Std Coef	Tolerance	t	P(2 Tail)
CONSTANT	0.102	0.064	0.000	.	1.600	0.110
A_PLAICE_DEL	0.677	0.014	0.914	1.000	49.408	0.000


Analysis of Variance

Source	Sum-of-Squares	df	Mean-Square	F-ratio	P
Regression	4602.279	1	4602.279	2441.151	0.000
Residual	904.940	480	1.885		


Case	86 is an outlier	(Studentized Residual =	4.381)
Case	88 has large leverage	(Leverage =	0.391)
Case	97 is an outlier	(Studentized Residual =	-8.786)
Case	229 has large leverage	(Leverage =	0.117)
Case	229 is an outlier	(Studentized Residual =	-9.536)
Case	233 has large leverage	(Leverage =	0.072)
Case	235 has large leverage	(Leverage =	0.067)
Case	248 is an outlier	(Studentized Residual =	4.296)
Case	360 is an outlier	(Studentized Residual =	-4.445)
Case	361 has large leverage	(Leverage =	0.111)
Case	361 is an outlier	(Studentized Residual =	12.285)
Case	481 has large leverage	(Leverage =	0.059)
Case	481 is an outlier	(Studentized Residual =	7.078)

Durbin-Watson D Statistic 2.125
First Order Autocorrelation -0.063

Plot of Residuals against Predicted Values



American plaice weights (kg)



Fourspot Flounder [cruise = Pooled data]

Dep Var: FOURSPOT_ALB N: 482 Multiple R: 0.932 Squared multiple R: 0.868

Adjusted squared multiple R: 0.868 Standard error of estimate: 0.861

Effect	Coefficient	Std Error	Std Coef	Tolerance	t	P(2 Tail)
CONSTANT	0.120	0.041	0.000	.	2.893	0.004
FOURSPOT_DEL	0.725	0.013	0.932	1.000	56.206	0.000


Analysis of Variance

Source	Sum-of-Squares	df	Mean-Square	F-ratio	P
Regression	2343.690	1	2343.690	3159.156	0.000
Residual	356.099	480	0.742		


Case	26 has large leverage	(Leverage =	0.037)
Case	53 has large leverage	(Leverage =	0.096)
Case	56 is an outlier	(Studentized Residual =	6.039)
Case	103 has large leverage	(Leverage =	0.056)
Case	103 is an outlier	(Studentized Residual =	6.099)
Case	145 has large leverage	(Leverage =	0.040)
Case	145 is an outlier	(Studentized Residual =	7.555)
Case	149 has large leverage	(Leverage =	0.096)
Case	155 has large leverage	(Leverage =	0.210)
Case	155 is an outlier	(Studentized Residual =	-6.757)
Case	157 has large leverage	(Leverage =	0.070)
Case	167 is an outlier	(Studentized Residual =	4.315)
Case	287 is an outlier	(Studentized Residual =	5.937)
Case	301 has large leverage	(Leverage =	0.047)
Case	305 is an outlier	(Studentized Residual =	-5.994)
Case	478 is an outlier	(Studentized Residual =	6.031)

Durbin-Watson D Statistic 2.127
First Order Autocorrelation -0.064

Plot of Residuals against Predicted Values



Fourspot flounder weights (kg)



Yellowtail Flounder [cruise = Pooled data]

Dep Var: YT_ALB N: 482 Multiple R: 0.941 Squared multiple R: 0.886

Adjusted squared multiple R: 0.886 Standard error of estimate: 1.887

Effect	Coefficient	Std Error	Std Coef Tolerance	t	P(2 Tail)
CONSTANT	0.146	0.088	0.000	1.655	0.099
YT_DEL	0.658	0.011	0.941	61.078	0.000


Analysis of Variance

Source	Sum-of-Squares	df	Mean-Square	F-ratio	P
Regression	13279.336	1	13279.336	3730.523	0.000
Residual	1708.629	480	3.560		


Case	21 has large leverage	(Leverage =	0.114)
Case	21 is an outlier	(Studentized Residual =	22.472)
Case	30 has large leverage	(Leverage =	0.279)
Case	31 has large leverage	(Leverage =	0.177)
Case	31 is an outlier	(Studentized Residual =	-8.421)
Case	92 is an outlier	(Studentized Residual =	-6.950)
Case	149 has large leverage	(Leverage =	0.194)
Case	151 is an outlier	(Studentized Residual =	5.007)
Case	174 is an outlier	(Studentized Residual =	5.075)

Durbin-Watson D Statistic 1.948
First Order Autocorrelation 0.023

Plot of Residuals against Predicted Values



Yellowtail flounder weights (kg)



Winter Flounder [cruise = Pooled data]

Dep Var: WINTER_F_ALB N: 482 Multiple R: 0.816 Squared multiple R: 0.666

Adjusted squared multiple R: 0.666 Standard error of estimate: 2.643

Effect	Coefficient	Std Error	Std Coef Tolerance	t	P(2 Tail)
CONSTANT	0.284	0.126	0.000	2.257	0.024
WINTER_F_DEL	0.777	0.025	0.816	30.956	0.000


Analysis of Variance

Source	Sum-of-Squares	df	Mean-Square	F-ratio	P
Regression	6691.408	1	6691.408	958.245	0.000
Residual	3351.831	480	6.983		


Case	81 has large leverage	(Leverage =	0.038)
Case	81 is an outlier	(Studentized Residual =	4.042)
Case	91 is an outlier	(Studentized Residual =	5.199)
Case	143 is an outlier	(Studentized Residual =	11.007)
Case	210 has large leverage	(Leverage =	0.068)
Case	210 is an outlier	(Studentized Residual =	5.565)
Case	239 has large leverage	(Leverage =	0.110)
Case	332 has large leverage	(Leverage =	0.147)
Case	332 is an outlier	(Studentized Residual =	-6.830)
Case	382 has large leverage	(Leverage =	0.042)
Case	382 is an outlier	(Studentized Residual =	4.839)
Case	383 has large leverage	(Leverage =	0.169)
Case	383 is an outlier	(Studentized Residual =	7.031)
Case	469 has large leverage	(Leverage =	0.116)
Case	469 is an outlier	(Studentized Residual =	-10.115)

Durbin-Watson D Statistic 1.893
First Order Autocorrelation 0.054

Plot of Residuals against Predicted Values



Winter flounder weights (kg)



Research Communications Branch
Northeast Fisheries Science Center
National Marine Fisheries Service, NOAA
166 Water St.
Woods Hole, MA 02543-1026

**MEDIA
MAIL**

Publications and Reports of the Northeast Fisheries Science Center

The mission of NOAA's National Marine Fisheries Service (NMFS) is "stewardship of living marine resources for the benefit of the nation through their science-based conservation and management and promotion of the health of their environment." As the research arm of the NMFS's Northeast Region, the Northeast Fisheries Science Center (NEFSC) supports the NMFS mission by "conducting ecosystem-based research and assessments of living marine resources, with a focus on the Northeast Shelf, to promote the recovery and long-term sustainability of these resources and to generate social and economic opportunities and benefits from their use." Results of NEFSC research are largely reported in primary scientific media (*e.g.*, anonymously-peer-reviewed scientific journals). However, to assist itself in providing data, information, and advice to its constituents, the NEFSC occasionally releases its results in its own media. Currently, there are three such media:

NOAA Technical Memorandum NMFS-NE -- This series is issued irregularly. The series typically includes: data reports of long-term field or lab studies of important species or habitats; synthesis reports for important species or habitats; annual reports of overall assessment or monitoring programs; manuals describing program-wide surveying or experimental techniques; literature surveys of important species or habitat topics; proceedings and collected papers of scientific meetings; and indexed and/or annotated bibliographies. All issues receive internal scientific review and most issues receive technical and copy editing.

Northeast Fisheries Science Center Reference Document -- This series is issued irregularly. The series typically includes: data reports on field and lab studies; progress reports on experiments, monitoring, and assessments; background papers for, collected abstracts of, and/or summary reports of scientific meetings; and simple bibliographies. Issues receive internal scientific review and most issues receive copy editing.

Resource Survey Report (formerly *Fishermen's Report*) -- This information report is a regularly-issued, quick-turnaround report on the distribution and relative abundance of selected living marine resources as derived from each of the NEFSC's periodic research vessel surveys of the Northeast's continental shelf. This report undergoes internal review, but receives no technical or copy editing.

TO OBTAIN A COPY of a *NOAA Technical Memorandum NMFS-NE* or a *Northeast Fisheries Science Center Reference Document*, either contact the NEFSC Editorial Office (166 Water St., Woods Hole, MA 02543-1026; 508-495-2350) or consult the NEFSC webpage on "Reports and Publications" (<http://www.nefsc.noaa.gov/nefsc/publications/>). To access *Resource Survey Report*, consult the Ecosystem Surveys Branch webpage (<http://www.nefsc.noaa.gov/femad/ecosurvey/mainpage/>).

ANY USE OF TRADE OR BRAND NAMES IN ANY NEFSC PUBLICATION OR REPORT DOES NOT IMPLY ENDORSEMENT.

Procedures for Issuing Manuscripts in the *Northeast Fisheries Science Center Reference Document (CRD) Series*

Clearance

All manuscripts submitted for issuance as CRDs must have cleared the NEFSC's manuscript/abstract/webpage review process. If any author is not a federal employee, he/she will be required to sign an "NEFSC Release-of-Copyright Form." If your manuscript includes material from another work which has been copyrighted, then you will need to work with the NEFSC's Editorial Office to arrange for permission to use that material by securing release signatures on the "NEFSC Use-of-Copyrighted-Work Permission Form."

For more information, NEFSC authors should see the NEFSC's online publication policy manual, "Manuscript/abstract/webpage preparation, review, and dissemination: NEFSC author's guide to policy, process, and procedure," located in the Publications/Manuscript Review section of the NEFSC intranet page.

Organization

Manuscripts must have an abstract and table of contents, and (if applicable) lists of figures and tables. As much as possible, use traditional scientific manuscript organization for sections: "Introduction," "Study Area" and/or "Experimental Apparatus," "Methods," "Results," "Discussion," "Conclusions," "Acknowledgments," and "Literature/References Cited."

Style

The CRD series is obligated to conform with the style contained in the current edition of the United States Government Printing Office Style Manual. That style manual is silent on many aspects of scientific manuscripts. The CRD series relies more on the CSE Style Manual. Manuscripts should be prepared to conform with these style manuals.

The CRD series uses the American Fisheries Society's guides to names of fishes, mollusks, and decapod

crustaceans, the Society for Marine Mammalogy's guide to names of marine mammals, the Biosciences Information Service's guide to serial title abbreviations, and the ISO's (International Standardization Organization) guide to statistical terms.

For in-text citation, use the name-date system. A special effort should be made to ensure that all necessary bibliographic information is included in the list of cited works. Personal communications must include date, full name, and full mailing address of the contact.

Preparation

Once your document has cleared the review process, the Editorial Office will contact you with publication needs – for example, revised text (if necessary) and separate digital figures and tables if they are embedded in the document. Materials may be submitted to the Editorial Office as files on zip disks or CDs, email attachments, or intranet downloads. Text files should be in Microsoft Word, tables may be in Word or Excel, and graphics files may be in a variety of formats (JPG, GIF, Excel, PowerPoint, etc.).

Production and Distribution

The Editorial Office will perform a copy-edit of the document and may request further revisions. The Editorial Office will develop the inside and outside front covers, the inside and outside back covers, and the title and bibliographic control pages of the document.

Once both the PDF (print) and Web versions of the CRD are ready, the Editorial Office will contact you to review both versions and submit corrections or changes before the document is posted online.

A number of organizations and individuals in the Northeast Region will be notified by e-mail of the availability of the document online.
